

Program Notice

FGIS PN-06-05

05-15-06

SAMPLE COLLECTION RESPONSIBILITIES FOR VERIFYING
THE ACCURACY OF MOISTURE METER CALIBRATIONS
CROP YEAR 2006

1. PURPOSE

This program notice transmits collection assignments for samples needed to verify the accuracy of official moisture meter calibrations. It also restates the procedure for collecting and submitting samples.

2. BACKGROUND

The annual Moisture Meter Calibration Study is conducted on current year crop samples to assess the accuracy of the official inspection system and of NTEP-certified moisture meters. FGIS moisture meter calibrations must be verified over the working moisture ranges, significant production areas, and relevant crop years. Each year, the evaluation is performed on samples submitted to the Quality Systems and Services Unit (QSS) from the field offices. After moisture testing, the samples are made available to other programs in the Technical Services Division.

Sample collection assignments for the respective offices are based on crop production within the geographic areas of responsibility. In some cases, additional assignments in the stable moisture ranges are given to export locations. Also, the quotas for corn, soybean and Hard Red Winter wheat are increased slightly to provide enough samples for the NTEP testing program.

It is understood that all requested moisture levels may not be available in all areas every year. Since a wide moisture range is very important to the study, field offices should make all reasonable efforts to provide the requested number of samples in each moisture range. However, extraordinary actions are not expected.

3. EFFECTIVE DATE

This program notice is effective upon receipt for the 2006 crop production. Wheat samples should be submitted by September 15, sunflower samples by November 15, and all other grain samples by November 1, 2006.

4. REPLACEMENT HIGHLIGHTS

This program notice supersedes FGIS PN 05-05, dated May 15, 2005.

5. RESPONSIBILITIES

The collection and submission of samples for the annual Moisture Meter Calibration Study are considered regular duties of the selected field offices. All associated time will be charged to the field office standardization management code.

6. ASSIGNMENTS FOR SELECTED FIELD OFFICES

During the 2006 growing season, the indicated numbers of samples of the commodities listed in Table 1 (Attachment 2) must be collected, tested for moisture, and submitted by the respective field offices to TSD-QSS. Each sample should weigh approximately 1500 grams.

7. INSTRUCTIONS

- a. The purpose of this effort is to obtain representative samples from the entire nation. Therefore, it is important to have each office fill its quota at all moisture levels, if possible. However, do not submit extra samples in any moisture range, and do not adjust the moisture level of samples by adding water or by drying in the laboratory.
- b. Samples with moisture levels slightly beyond the established moisture ranges are useful in calibrating the extreme ends of the calibrations and extending the measurement ranges. For this reason, the ranges of requested samples (Table 1) have been extended slightly beyond established limits. When submitting samples, if the moisture falls outside the range of the applicable GAC 2100 calibration, obtain an approximate moisture. The true moisture will later be determined at TSD by air oven.
- c. If dockage is removed for inspection purposes, do not recombine it before submitting the sample.
- d. The significant amount of time and effort invested in collecting and submitting the moisture samples can easily be lost through insect damage, microbial spoilage, or late sample submission. To prevent such loss, please collect the samples during the growing season and at harvest time and submit them promptly.

Samples above 16% (above 14% for sunflower seeds and 11% for minor oilseeds) require special handling. To minimize loss by spoilage, keep high moisture samples refrigerated (not frozen) until shipped and ship the samples by Federal Express at least 48 hours before a weekend/holiday.

- e. An easy way to account for samples submitted is to prepare mailing tags for the total number of samples of each commodity to be collected. On the back of each tag, write the commodity and moisture range. When all of the mailing tags are used, the collection assignment has been met.
- f. Most dielectric moisture meters have a built-in test weight correction. These corrections need to be checked using external test weight data. For samples of sufficient volume, test weight will be determined by TSD-QSS, so it is not necessary to record test weight on the mailing tag. However, some submitted samples are too small to fill the kettle. For such samples, please record the test weight on the tag (or transmittal slip) if it is known.
- g. Questions concerning these instructions should be directed to Patricia Jackson (816) 891-0450. If there is a special problem with a sample assignment, please notify the Moisture Laboratory, TSD-QSS, by telephone as early in the season as possible.
- h. Seal each sample in a polyethylene bag (6 mil thickness) and insert the bag into a canvas grain bag. When shipping several samples in a larger container (box or mail sack), a canvas grain bag around each poly bag will help prevent the poly bags from breaking in transit. Record the field office location, date, commodity, official meter moisture, and test weight (if sample size is limited) on the back of the mailing tag or transmittal form [Attachment 1] accompanying the sample. Attach the mailing tag to the bag. Send samples to:

USDA-GIPSA-FGIS Technical Center
 Technical Services Division
 Moisture Laboratory
 10383 N. Ambassador Drive
 Kansas City, MO 64153-1394

/s/ John Giler

John Giler, Acting Director
 Field Management Division
 Attachments

Moisture Sample Transmittal Form

Field Office Use Only:

OFFICE _____ MOISTURE _____

DATE _____ TEST WT. _____

COMMODITY _____

ISE Use Only: *Date Received*

Moisture Sample Transmittal Form

Field Office Use Only:

OFFICE _____ MOISTURE _____

DATE _____ TEST WT. _____

COMMODITY _____

ISE Use Only: *Date Received*

Moisture Sample Transmittal Form

Field Office Use Only:

OFFICE _____ MOISTURE _____

DATE _____ TEST WT. _____

COMMODITY _____

ISE Use Only: *Date Received*

Moisture Sample Transmittal Form

Field Office Use Only:

OFFICE _____ MOISTURE _____

DATE _____ TEST WT. _____

COMMODITY _____

ISE Use Only: *Date Received*

Moisture Sample Transmittal Form

Field Office Use Only:

OFFICE _____ MOISTURE _____

DATE _____ TEST WT. _____

COMMODITY _____

ISE Use Only: *Date Received*

Moisture Sample Transmittal Form

Field Office Use Only:

OFFICE _____ MOISTURE _____

DATE _____ TEST WT. _____

COMMODITY _____

ISE Use Only: *Date Received*

Table 1. Sample collection assignments, 2006 Crop Year

1. Barley, Six-Rowed	Office	Moisture Range (%)						
		<u>7-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>		
	California	2	2	2	1	7		
	Grand Forks	5	6	6	4	21		
	Minneapolis	6	7	7	4	24		
	Moscow	2	2	2	1	7		
	Toledo	2	4	3	2	11		
2. Barley, Two-Rowed	Office	Moisture Range (%)						
		<u>7-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>		
	Grand Forks	2	2	2	1	7		
	Minneapolis	2	2	2	1	7		
	Moscow	10	11	10	6	37		
	Washington	5	7	4	3	19		
3. Corn	Office	Moisture Range (%)						
		<u>7-11</u>	<u>11-14</u>	<u>14-18</u>	<u>18-22</u>	<u>22-26</u>	<u>26-31</u>	<u>All</u>
	Cedar Rapids	12	14	16	11	10	10	73
	Grand Forks	2	2	2	2	1	1	10
	League City	1	3	3	2	2	1	12
	Minneapolis	8	9	11	9	7	6	50
	New Orleans	1	2	2	0	0	0	5
	Stuttgart	3	5	4	4	2	2	20
	Toledo	8	9	12	9	8	6	52
	Wichita	11	13	14	13	12	10	73
4. Oats	Office	Moisture Range (%)						
		<u>7-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>		
	Cedar Rapids	7	9	9	3	28		
	Grand Forks	2	4	3	2	11		
	Minneapolis	10	11	9	6	36		
	Wichita	1	2	2	0	5		

5. Rough Rice, Long Grain	Office	Moisture Range (%)					<u>All</u>
		<u>7-11</u>	<u>11-14</u>	<u>14-18</u>	<u>18-22</u>	<u>22-26</u>	
	League City	3	4	4	3	2	16
	New Orleans	6	6	6	6	5	29
	Stuttgart	9	10	9	9	8	45

6. Rough Rice, Medium Grain	Office	Moisture Range (%)					<u>All</u>
		<u>7-11</u>	<u>11-14</u>	<u>14-18</u>	<u>18-22</u>	<u>22-26</u>	
	California	10	12	12	11	9	54
	New Orleans	2	2	2	1	1	8
	Stuttgart	6	7	6	5	4	28

7. Sorghum	Office	Moisture Range (%)					<u>All</u>
		<u>7-11</u>	<u>11-14</u>	<u>14-18</u>	<u>18-22</u>	<u>22-26</u>	
	League City	4	5	4	3	2	18
	New Orleans	2	3	3	0	0	8
	Stuttgart	2	3	4	0	0	9
	Wichita	10	10	9	7	4	40

8. Soybeans	Office	Moisture Range (%)					<u>All</u>
		<u>7-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>		
	Cedar Rapids	13	14	13	9	49	
	Grand Forks	3	4	3	2	12	
	Minneapolis	10	10	9	8	37	
	New Orleans	4	5	2	2	13	
	Stuttgart	7	7	6	5	25	
	Toledo	10	11	10	8	39	
	Wichita	12	13	11	9	45	

9. Sunflower Seed, Oil Type	Office	Moisture Range (%)						<u>All</u>
		<u>4-7</u>	<u>7-10</u>	<u>10-14</u>	<u>14-18</u>	<u>18-22</u>	<u>22-26</u>	
	Grand Forks	6	8	7	7	6	5	39
	Minneapolis	10	12	11	9	8	7	57
	Wichita	5	7	6	6	6	4	34

10. Wheat, Durum	Office	Moisture Range (%)				
		<u>6-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>
	California	3	3	3	1	10
	Duluth	2	2	2	0	6
	Grand Forks	4	6	5	3	18
	Minneapolis	3	4	4	2	13
	Moscow	3	4	4	2	13
	Wichita	3	3	3	1	10

11. Wheat, Hard Red Spring	Office	Moisture Range (%)				
		<u>6-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>
	Duluth	2	2	2	0	6
	Grand Forks	5	7	6	4	22
	Minneapolis	7	9	7	5	28
	Moscow	4	5	5	2	16
	Washington	1	2	1	0	4

12. Wheat, Hard Red Winter	Office	Moisture Range (%)				
		<u>7-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>
	California	3	3	2	1	9
	League City	3	3	2	1	9
	Minneapolis	4	4	4	1	13
	Moscow	4	4	4	2	14
	Wichita	12	13	12	8	45

13. Wheat, Hard White	Office	Moisture Range (%)				
		<u>6-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>
	California	4	4	3	2	13
	Moscow	6	7	7	5	25
	Washington	4	4	4	2	14
	Wichita	5	6	5	2	18

14. Wheat, Soft Red Winter	Office	Moisture Range (%)				
		<u>6-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>
	Cedar Rapids	3	4	3	2	12
	New Orleans	2	3	2	1	8
	Stuttgart	4	4	3	2	13
	Toledo	5	7	6	4	22
	Wichita	3	5	4	3	15

15. Wheat, Soft White	Office	Moisture Range (%)				
		<u>7-11</u>	<u>11-14</u>	<u>14-17</u>	<u>17-21</u>	<u>All</u>
	Moscow	7	9	7	4	27
	Toledo	3	4	3	1	11
	Washington	8	9	9	6	32

The majority of the following requested samples should reflect typical market moisture ranges. When available, one or two samples of each grain or commodity, outside the typical range, would be beneficial to the moisture calibration program. The samples should represent diverse growing conditions.

16. Beans, Black	Office	<u>Number of Samples</u>
	Grand Forks	5
	Minneapolis	5
	Moscow	5
	Toledo	5

17. Beans, Blackeye	Office	<u>Number of Samples</u>
	California	10
	Wichita	10

18. Beans, Cranberry	Office	<u>Number of Samples</u>
	California	10
	Minneapolis	10
	Toledo	10

19. Beans, Garbanzo	Office	<u>Number of Samples</u>
	California	5
	Grand Forks	5
	Minneapolis	5
	Moscow	5
	Washington	5

20. Beans, Great Northern	Office	<u>Number of Samples</u>
	Moscow	10
	Wichita	10

21. Beans, Kidney (Light and Dark)	Office	<u>Number of Samples</u>
	California	5
	Grand Forks	5
	Minneapolis	5
	Toledo	5
	Wichita	5

22. Beans, Baby Lima	Office	<u>Number of Samples</u>
	California	10

23. Beans, Large Lima	Office	<u>Number of Samples</u>
	California	10

24. Beans, Pea (Navy)	Office	<u>Number of Samples</u>
	Grand Forks	10
	Minneapolis	10
	Toledo	10

25. Beans, Pink	Office	<u>Number of Samples</u>
	California	5
	Duluth	5
	Grand Forks	5
	Minneapolis	5
	Moscow	5
	Washington	5

26. Beans, Pinto	Office	<u>Number of Samples</u>
	California	5
	Grand Forks	5
	Minneapolis	5
	Moscow	5
	Wichita	5

27. Beans, Small Red	Office	<u>Number of Samples</u>
	Grand Forks	5
	Minneapolis	5
	Moscow	5
	Toledo	5
	Washington	5

28. Canola	Office	<u>Number of Samples</u>
	Duluth	5
	Grand Forks	5
	Minneapolis	5
	Washington	5

29. Flaxseed	Office	<u>Number of Samples</u>
	Grand Forks	10
	Minneapolis	10

30. Lentils	Office	<u>Number of Samples</u>
	Moscow	5
	Grand Forks	5
	Minneapolis	5
	Washington	5
31. Mustard Seed, Yellow	Office	<u>Number of Samples</u>
	Grand Forks	10
	Moscow	10
	Washington	10
32. Peas, Smooth Dry	Office	<u>Number of Samples</u>
	Grand Forks	10
	Moscow	10
	Washington	10
33. Rice, Long Grain Brown	Office	<u>Number of Samples</u>
	League City	10
	New Orleans	10
	Stuttgart	10
34. Rice, Long Grain Milled	Office	<u>Number of Samples</u>
	League City	10
	New Orleans	10
	Stuttgart	10
35. Rice, Medium Grain Brown	Office	<u>Number of Samples</u>
	League City	5
	New Orleans	5
	California	10
	Stuttgart	5

36. Rice, Medium Grain Milled	Office	<u>Number of Samples</u>
	League City	5
	New Orleans	5
	California	10
	Stuttgart	5

37. Rice, Short Grain Rough	Office	<u>Number of Samples</u>
	California	10

38. Rye	Office	<u>Number of Samples</u>
	Grand Forks	5
	Minneapolis	5
	Toledo	5
	Wichita	5

39. Safflower Seed	Office	<u>Number of Samples</u>
	California	5
	Grand Forks	5
	Minneapolis	5
	Moscow	5
	Wichita	5

40. Sunflower Seed, Confectionary	Office	<u>Number of Samples</u>
	Grand Forks	10
	Minneapolis	10
	Wichita	10